

**FP2020  
2015-2016 PROGRESS REPORT**

**INDICATOR NO. 14  
ESTIMATE TABLE**

COUNTRY	METHOD INFORMATION INDEX	PERMANENT	LONG-ACTING		SHORT-TERM	
	TOTAL	STERILIZATION (FEMALE)	IUD	IMPLANT	INJECTABLE	PILL
Burkina Faso	40.5%			48.8%	40.9%	43.5%
Cambodia	67.4%	62.5%	86.7%	80.4%	70.1%	59.2%
Comoros	36.2%			51.4%	30.0%	40.1%
DR Congo	28.4%	6.4%		50.4%	35.6%	12.0%
Egypt	28.8%	25.4%	30.4%	27.3%	30.0%	25.5%
Ethiopia	39.6%	41.4%*	60.2%	47.0%	36.5%	50.3%
Gambia	31.0%		29.9%*		33.5%	26.5%
Ghana	43.7%	35.7%*	61.6%*	62.4%	60.7%	29.7%
Guinea	31.3%			0.0%	29.1%	28.6%
Haiti	51.7%	30.0%		62.1%	54.3%	38.7%
Indonesia	30.4%	27.3%	39.7%	34.1%	31.2%	32.5%
Kenya	43.2%	36.2%	63.2%	54.7%	38.9%	34.3%
Kyrgyzstan	56.2%	26.9%*	59.5%	0.0%	0.0%	46.5%
Lesotho	27.0%	3.9%*	55.5%*	44.3%	26.7%	24.4%
Liberia	61.4%			75.6%	62.3%	51.9%
Mali	33.3%		50.1%*	41.8%	31.2%	25.4%
Niger	28.4%		0.0%	39.8%*	34.4%	25.5%
Nigeria	47.1%	23.8%*	64.4%	73.3%	52.5%	30.9%
Pakistan	13.5%	7.6%	20.6%		18.3%	11.2%
Philippines	52.1%	45.9%	69.5%		58.5%	50.5%
Rwanda	57.9%	23.9%	65.1%	61.4%	59.1%	53.0%
Senegal	64.8%		72.4%*	54.4%	74.2%	63.3%
Sierra Leone	69.8%	54.0%*	76.7%	78.0%	72.1%	59.7%
Tajikistan	59.4%		59.1%	0.0%	65.4%	60.7%
Togo	67.5%		72.0%	79.8%	68.8%	44.4%
Uganda	48.3%	46.9%		78.3%	45.7%	50.6%
Yemen	34.9%	22.0%	45.7%	44.8%	36.3%	30.9%
Zambia	71.8%	49.9%	82.3%	83.8%	73.8%	62.4%

\*Small sample size (between 25 and 50).

\*\*Among all women who responded to this set of three questions, not just among those who were told about side effects.

**Note:** Blank cells indicate that the sample size was too small for inclusion or no data was available. No countries had a large enough sample size to analyze male sterilization.

**DEFINITION:**

An index measuring the extent to which women were given specific information when they received family planning services. The index is composed of three questions: 1) Were you informed about other methods? 2) Were you informed about side effects? 3) Were you told what to do if you experienced side effects? The reported Method Information Index value is the percent of women who responded “yes” to all three questions.

**SOURCE:**

For each country, the most recent national survey (DHS, PMA2020). Data reflect all women, except Egypt, Pakistan and Yemen, which reflect married or in-union women.

RESPONSES TO INDIVIDUAL QUESTIONS AMONG USERS OF MODERN METHODS				
COUNTRY	TOLD OF OTHER METHODS	TOLD ABOUT SIDE EFFECTS	TOLD WHAT TO DO ABOUT SIDE EFFECTS**	SOURCE
Burkina Faso	71.1%	58.3%	48.9%	2015 PMA2020 R2
Cambodia	75.7%	79.6%	77.5%	2014 DHS
Comoros	62.2%	54.5%	45.7%	2012 DHS
DR Congo	50.8%	57.2%	47.5%	2013-14 DHS
Egypt	56.0%	45.0%	34.5%	2014 DHS
Ethiopia	62.2%	52.5%	43.3%	2015 PMA2020 R3
Gambia	57.5%	47.2%	41.8%	2013 DHS
Ghana	65.1%	58.5%	49.2%	2015 PMA2020 R4
Guinea	48.6%	48.6%	43.1%	2012 DHS
Haiti	64.6%	70.2%	63.7%	2012 DHS
Indonesia	57.6%	49.2%	36.8%	2015 PMA2020 R1
Kenya	70.4%	54.5%	51.9%	2014 DHS
Kyrgyzstan	64.6%	70.5%	67.1%	2012 DHS
Lesotho	62.6%	39.8%	36.5%	2014 DHS
Liberia	72.0%	75.0%	72.9%	2013 DHS
Mali	56.8%	53.1%	46.3%	2012-13 DHS
Niger	55.9%	39.6%	35.4%	2012 DHS
Nigeria	64.8%	60.3%	54.8%	2013 DHS
Pakistan	28.2%	34.0%	28.1%	2012-13 DHS
Philippines	71.4%	67.8%	67.9%	2013 DHS
Rwanda	79.5%	64.8%	68.5%	2014-15 DHS
Senegal	84.5%	72.7%	76.8%	2014 DHS
Sierra Leone	82.7%	75.7%	74.9%	2013 DHS
Tajikistan	68.1%	77.0%	71.8%	2012 DHS
Togo	82.7%	78.1%	74.6%	2013-14 DHS
Uganda	64.2%	62.1%	54.2%	2015 PMA2020 R3
Yemen	57.0%	55.7%	45.8%	2013 DHS
Zambia	83.3%	79.7%	78.1%	2013-14 DHS

\*Small sample size (between 25 and 50).

\*\*Among all women who responded to this set of three questions, not just among those who were told about side effects.

**Note:** Blank cells indicate that the sample size was too small for inclusion or no data was available. No countries had a large enough sample size to analyze male sterilization.